

Intestiphage® - Patient leaflet (Instruction for medical use)

The product contains bacteriophages killing the pathogenic and opportunistic pathogenic bacteria in the intestine. The product contains bacteriophages specifically affection the following bacteria:

Shigella flexneri
Shigella sonnei
Salmonella enterica
Escherichia coli
Pseudomonas aeruginosa
Proteus vulgaris
Proteus mirabilis.
Enterococcus faecalis
Staphylococcus aureus

The bacteriophages contained in the product proliferate inside the bacterial cells and destroy them. The bacteriophages also induce activation of immune reactions in the body, preventing recurrence of infectious disease and transition of the inflammatory process into a chronic form.

The product does not violate the condition of the normal microflora in the gastro-intestinal tract.

Intestiphage®

The full information on the product is given in the instruction for medical use. Before use read the Instructions for medical use of Intestiphage® medicinal product.

Indications for use of Intestiphage® medicinal product

- Salmonellosis;
- Shigellosis;
- Dysbacteriosis;
- Infectious colitis and enterocolitis;

Drug form

Solution in vial.

Administration of Intestiphage® medicinal product

The treatment of pyoinflammatory diseases with a localized size of disease is carried out locally as well as orally.

For treatment the bacteriophage is administered orally 4 times a day 1 hour before meals.

Rectally (in enemas) the product is administered 1 time a day after defecation instead of 1 oral administration.

The duration of treatment is 7-10 days.

Dosage

The dosage of product depends on the age of the patient and the way of administration.

Age	Dosage per 1 administration (ml)	
	orally	rectally (enemas)
from 2 to 6 month	5	10
from 6 to 12 month	10	20
from 1 to 3 years	15	30
from 3 to 8 years	20	40
8 years and older	30	50

As a preventive measure bacteriophage is administered orally according to the age dose 1 time a day 1 hour before meals. The duration of the course depends on the epidemic situation.

Side effects of Intestiphage® medicinal product

No cases of adverse reactions were described.

Contraindications to administration of Intestiphage® medicinal product

Hypersensitisation to the product components.

Administration of Intestiphage® medicinal product during pregnancy and breastfeeding period

The safety and efficiency of the product during pregnancy and breastfeeding period is not yet ascertained, therefore it is advisable to consult your physician before use.

Drug-to-drug interaction

The product can be used along with the other antibacterial agents.

Administration details of Intestiphage® medicinal product

The essential condition for an efficient phagotherapy is a prior definition of the causative bacterium phagosensitivity. The product should be administered as soon as possible.

If you use only a part of the solution from the vial per administration, the following rules should be observed:

- thoroughly wash your hands before opening the vial,
- take off the cap without removing the stopper,
- withdraw the solution from the vial only using a sterile syringe by the way of puncturing the stopper,
- don't leave the vial opened, keep in the fridge.

Caution

The information on the product is presented for information purposes only and shall not be used as a guide for self-treatment. Only a doctor can make a decision as to prescribing of the product, as well as doses thereof and method of administration.

Manufacturer

Intestiphage® medicinal product is manufactured by "PHARMEX GROUP" LLC, Ukraine, for NeoProbioCare Inc., Canada.